

*Short-Communication***The Role of Urban Forests as Carbon Sink: A Case Study in the Urban Forest of Banda Aceh, Indonesia**Ali Muhammad Muslih<sup>1,\*</sup>, Asyrafun Nisa<sup>1</sup>, Sugianto Sugianto<sup>2</sup>, Tuti Arlita<sup>1</sup>, Subhan<sup>1</sup><sup>1</sup> Department of Forestry, Faculty of Agriculture, Universitas Syiah Kuala. Jl. Tengku Hasan Krueng Kale No. 3, Kopelma Darussalam, 23111, Banda Aceh, Indonesia<sup>2</sup> Department of Soil Science, Faculty of Agriculture, Universitas Syiah Kuala. Jl. Tengku Hasan Krueng Kale No. 3, Kopelma Darussalam, 23111, Banda Aceh, Indonesia\* Corresponding Author. E-mail address: [alimmuslih@unsyiah.ac.id](mailto:alimmuslih@unsyiah.ac.id)

## ARTICLE HISTORY:

Received: 22 May 2022

Peer review completed: 18 June 2022

Received in revised form: 27 June 2022

Accepted: 17 August 2022

## KEYWORDS:

*Carbon sequestration**Carbon stock**Green open space**Tree biomass**Urban forest*

## ABSTRACT

Forests have a role as carbon sinks and come under the spotlight when the world is facing the climate change problem. Urban forests have functions as the most effective carbon sinks. Banda Aceh City only has 7.15 ha of urban forest, and the vegetation that grows in the urban forest is essential for carbon emissions reduction. This study aimed to determine the potential tree biomass and carbon sink value in the BNI Urban Forest, Banda Aceh City, Indonesia. This study used vegetation analysis by inventory method with the census sampling method. The diameter of all trees was measured, and tree biomass, carbon stocks, and carbon sequestration were calculated. This study identified 16 tree species in the BNI Urban Forest with a total of 318 individuals, showing *Casuarina equisetifolia* as the most dominant tree (163 trees). The above-ground biomass, carbon stock, and carbon sequestration by the vegetation in the BNI Urban Forest were 24.66 tons/ha, 11.59 tons/ha, and 42.51 tons/ha, respectively. This study presented the vegetation conditions and the potential of carbon sequestration, which can be used to monitor and evaluate the benefits of the existence of urban forests in Banda Aceh.

© 2022 The Author(s). Published by Department of Forestry, Faculty of Agriculture, the University of Lampung in collaboration with Indonesia Network for Agroforestry Education (INAFE). This is an open-access article under the CC BY-NC license: <https://creativecommons.org/licenses/by-nc/4.0/>.

**1. Introduction**

Forests have a significant role in the survival and life of living things. Therefore, forest management is essential. It is helpful to know the extent of utilization and use of forest areas (De Groot et al. 2019). The following points are strongly related to the application of forest ecosystem service functions that will have a good impact on the environment: (1) forests have a role as a carbon sink and begin to be in the spotlight when the earth is faced with the problem of global climate change caused (Subhan et al. 2021; Wahanisa 2015), (2) awareness of the importance of protecting and preserving the environment has increased along with global climate change, which has become a world issue in recent years (Hyde et al. 2022), and (3) global warming, caused by large quantities of carbon dioxide (CO<sub>2</sub>) gas in the atmosphere (Skytt et al. 2020), triggering an increase in the earth's temperature due to the current greenhouse effect (Xu and Cui 2021).

The effort to reduce greenhouse gas concentrations (emissions) in the atmosphere is to reduce the release of CO<sub>2</sub> into the air (Fenton 2017). For this reason, the amount of CO<sub>2</sub> in the air must be controlled by increasing the amount of CO<sub>2</sub> uptake by plants as much as possible and

suppressing the release of greenhouse gas as low as possible. Naturally, the release of CO<sub>2</sub> into the atmosphere occurs through various mechanisms such as the respiration of living things, decomposition of organic matter, and combustion of biomass (Marchi et al. 2016). In addition, plants absorb CO<sub>2</sub> during photosynthesis to convert CO<sub>2</sub> into oxygen (O<sub>2</sub>) (Sharkey 2021).


Banda Aceh City has an urban forest named BNI Urban Forest that was initiated after the tsunami in 2004 with a large existing urban forest of about 7.15 ha (Alfida et al. 2016). Urban forests are part of green open spaces, which function as the most effective carbon sinks and can reduce greenhouse gas emissions in the atmosphere (Lee et al. 2021; Mahmoudi et al. 2022). Besides adding to the city's beauty, the existence of urban forests can also absorb CO<sub>2</sub> in the air through the photosynthesis process, which can contribute to climate change mitigation (Devisscher et al. 2022; Mukhlison 2013). The different types of trees will have different abilities to absorb CO<sub>2</sub> in the air. Therefore, the selection of species that meet the criteria for selecting urban forest types (Conway and Vecht 2015; Morakinyo et al. 2020; Sun et al. 2022; Wu et al. 2021) and information on the amount of carbon stored by these types is important for urban forest planners to plan the development of urban forests and to absorb a high amount of carbon (Barona et al. 2022; Petri et al. 2019).

Banda Aceh City only has one urban forest as green open spaces and it has an essential role as a carbon sink. However, no previous study on carbon stock estimation and carbon sequestration was conducted on this site. Therefore, this study aimed to determine the potential carbon stocks and carbon sequestration in the BNI Urban Forest, Banda Aceh.

## 2. Materials and Methods

### 2.1. Study Area

This research was carried out from July 2021 to October 2021 in the BNI Urban Forest, Banda Aceh City (5.584585, 95.351010), with a total area of 7.15 ha. The research location was in Tibang Village, Syiah Kuala District, Banda Aceh City, with a study area of 5.48 ha (**Fig. 1**).



**Fig 1.** Research location in BNI Urban Forest, Banda Aceh, Indonesia.

## 2.2. Materials

The tools used were a global positioning system (GPS) tracker, roll meters (50 m), phiband for measuring tree diameter at breast height (DBH), camera, and tally sheet. The objects observed were all the trees in BNI Urban Forest Banda Aceh.

## 2.3. Methods

The research was conducted utilizing the vegetation analysis inventory method (Muslih et al. 2021; Ng et al. 2021). In addition, the census sampling method was carried out by recording all species and numbers of trees above ground.

### 2.3.1. Data collection

The data of all species and numbers of trees in the BNI Urban Forest Banda Aceh were collected. Trees that became the object of this study were trees having a DBH of more than 20 cm. Data collection was carried out using the vegetation inventory method (Rahmadwiati et al. 2022) and identifying plant diversity (Sugianto et al. 2021).

### 2.3.2. Data analysis

The data from the field was then calculated, including the tree diameter, tree biomass, carbon stocks, and carbon sequestration.

#### 2.3.2.1. Tree biomass

Tree biomass is determined based on the allometric formula for tropical wood species in Indonesia (Krisnawati et al. 2012). The allometric used is allometric for mixed trees because tree stands in urban forests are highly varied.

$$AGB = -2,75 + 2,591 * \ln(D) \quad (1)$$

where  $AGB$  is above-ground biomass (kg), and  $D$  is tree diameter.

#### 2.3.2.2. Carbon stocks potential

After obtaining the biomass amount, the carbon content of the biomass was calculated using Equation 2 (Badan Standardisasi Nasional 2011):

$$Cb = AGB \times \% C \text{ organic} \quad (2)$$

where  $Cb$  is the carbon content of biomass (kg),  $AGB$  is above-ground biomass (kg), and  $\% C \text{ organic}$  is a percentage value of carbon content (0.47).

The carbon stock was then calculated using Equation 3 (Badan Standardisasi Nasional 2011):

$$C = \frac{\sum C_b}{A} \quad (3)$$

where  $C$  is potential carbon stock (ton/ha),  $C_b$  is carbon content of biomass (ton), and  $A$  is research area (ha).

### 2.3.2.3. Carbon sequestration

Carbon sequestration was calculated by multiplying the carbon content of biomass by the carbon conversion factor (Ali et al. 2022; Tooichi 2018).

$$CO_2 = Cb \times 3.67 \quad (4)$$

where  $CO_2$  is carbon sequestration,  $Cb$  is the carbon content of biomass (ton/ha), and 3.67 is the conversion of element C to  $CO_2$ .

## 3. Results and Discussion

### 3.1. Tree Composition

The BNI Urban Forest Banda Aceh previously was only empty land caused by the impact of the tsunami in 2004. In 2011, the government developed the area into a cleaner and more organized urban forest as one of the green open spaces that continues to change until now. Currently, the benefits of the BNI Urban Forest Banda Aceh have been felt by the people of Banda Aceh City with the available biodiversity, improved water and soil quality, and the BNI Urban Forest Banda Aceh which is now greener and more beautiful. So that it can be said that the existence of trees that grow there enters into plants a value that can be felt (Muslih et al. 2022). Subhan et al. (2022) explained that the existence of a well-maintained urban forest would become a habitat for animals such as primates, reptiles, and birds.

Trees grow in groups consisting of 16 tree species with a total of 318 individuals. The BNI Urban Forest Banda Aceh area is dominated by *Casuarina equisetifolia*. The composition of species and number of trees in the BNI Banda Aceh can be seen in **Table 1**.

**Table 1.** Species composition and number of trees in BNI Urban Forest Banda Aceh

No.	Species	Family	Number of tree (individual)	Percentage (%)
1	<i>Acacia mangium</i>	Mimosaceae	6	1.9
2	<i>Pterocarpus indicus</i>	Fabaceae	15	4.7
3	<i>Tamarindus indica</i>	Fabaceae	13	4.1
4	<i>Pithecellobium dulce</i>	Fabaceae	7	2.2
5	<i>Ficus benjamina</i>	Moraceae	3	0.9
6	<i>Casuarina equisetifolia</i>	Casuarinaceae	163	51.3
7	<i>Delonix regia</i>	Fabaceae	2	0.6
8	<i>Senna siamea</i>	Fabaceae	5	1.6
9	<i>Lannea grandis</i>	Anacardiaceae	2	0.6
10	<i>Sterculia foetida</i>	Malvaceae	22	6.9
11	<i>Swietenia mahagoni</i>	Meliaceae	3	0.9
12	<i>Azadirachta indica</i>	Meliaceae	1	0.3
13	<i>Hura crepitans</i>	Euphorbiaceae	8	2.5
14	<i>Albizia chinensis</i>	Fabaceae	3	0.9
15	<i>Samanea saman</i>	Fabaceae	60	18.9
16	<i>Hibiscus tiliaceus</i>	Malvaceae	5	1.6
	Total		318	100

The most dominant tree was *C. equisetifolia* with 163 individuals (51.3%). This species is the dominant tree because BNI Urban Forest Banda Aceh is located in a coastal area where sea pine has good growth potential on nutrient-poor land. The least number is *Azadirachta indica*, with

only 1 individual (0.3%). Based on the Republic of Indonesia Government Regulation Number 63 of 2002, one of the urban forests' functions is to improve and maintain the microclimate and aesthetic value following its function. Based on its topographical location in coastal areas, many *C. equisetifolia* trees are found. *C. equisetifolia* has a function in efforts to improve the microclimate, therefore improving the lighting conditions of sunlight, air pressure, and relative humidity (Harjadi 2017).

The second most dominant species in the BNI Urban Forest Banda Aceh is *Samanea saman*. This tree strongly supports the need for shade trees and producers of fresh urban air for visitors who carry out activities in the area. According to Sumardi and Widyastuti (2013), the urban forest needs trees that grow large and shady to support social functions, environmental conservation, and aesthetics. It is the reason shady trees are more commonly found in urban forests.

### 3.2. Tree Biomass and Carbon Stock

**Table 2** shows that the amount of biomass in the Banda Aceh City Forest area is 24,669.18 kg/ha. In addition, the potential carbon stock is 11,594.51 kg/ha, and CO<sub>2</sub> sequestration is 42.51 tons/ha. The largest tree biomass and carbon stock were found in *C. equisetifolia*, while the smallest tree biomass and carbon stock were found in *A. indica*. This number will continue to increase along with the growth and development of trees (Ledheng et al. 2022) because the vegetation in the BNI Urban Forest Banda Aceh is still under maintenance.

**Table 2.** The result analysis of biomass, carbon stocks, and carbon sequestration

No.	Species	Number of Tree (individual)	AGB (kg)	C (kg)	CO <sub>2</sub> (ton)
1	<i>Acacia mangium</i>	6	1,991.65	936.08	3.43
2	<i>Pterocarpus indicus</i>	15	5,246.78	2,465.99	9.04
3	<i>Tamarindus indica</i>	13	4,006.77	1,883.18	6.91
4	<i>Pithecellobium dulce</i>	7	1,945.35	914.31	3.35
5	<i>Ficus benjamina</i>	3	1,307.89	614.71	2.25
6	<i>Casuarina equisetifolia</i>	163	67,717.95	31,827.43	116.70
7	<i>Delonix regia</i>	2	392.15	184.31	0.68
8	<i>Senna siamea</i>	5	2,809.63	1,320.53	4.84
9	<i>Lannea grandis</i>	2	682.19	320.63	1.18
10	<i>Sterculia foetida</i>	22	7,348.12	3,453.62	12.66
11	<i>Swietenia mahagoni</i>	3	571.15	268.44	0.98
12	<i>Azadirachta indica</i>	1	229.48	107.86	0.40
13	<i>Hura crepitans</i>	8	2,739.90	1,287.75	4.72
14	<i>Albizia chinensis</i>	3	2,970.23	1,396.01	5.12
15	<i>Samanea saman</i>	60	33,423.34	15,708.97	57.60
16	<i>Hibiscus tiliaceus</i>	5	1,804.55	848.14	3.11
Total		318	135,187.12	63,537.95	232.97
kg/ha		-	24,669.18	11,594.51	-
Ton/ha		-	24.66	11.59	42.51

Notes: AGB is above-ground biomass, C is potential carbon stock, and CO<sub>2</sub> is carbon sequestration.

Several studies related to the estimation of carbon stocks have been carried out. For example, Baruna et al. (2017) conducted research in the Urban Forest of the Pontianak City Hall and reported a carbon stock of 70.27 tons/ha. It can be seen that the carbon stocks in the BNI Urban Forest Banda Aceh are very small when compared to the carbon stock in the Urban Forest of the Pontianak City Hall. The different characteristics of each region cause these differences. Several

factors can influence genetic factors, location, soil conditions, stand density, and management practices applied by each manager.

The biomass content and potential carbon stocks contained in the BNI Urban Forest have different amounts. An unbalanced tree vegetation structure causes this, ranging from tree species, where they grow, cropping forms still piling up, and uneven planting locations. Reducing the number of trees per hectare does not reduce the amount of carbon sequestration per hectare (Matthews et al. 2020). It was due to a significant increase in stem diameter, some leaves, and some stomata. These differences significantly affect the conditions of CO<sub>2</sub> sequestration and stored carbon content.

### 3.3. Carbon Sequestration

The potential CO<sub>2</sub> sequestration in BNI Urban Forest Banda Aceh was 232.97 tons, or 42.51 tons/ha (Table 2). It can be seen that the largest CO<sub>2</sub> uptake of 116.70 tons was found in *C. equisetifolia*, with a total of 163 trees. Plants at the tree growth stage will produce a higher amount of biomass and carbon than plants at the sapling and pole stages. The results show that the amount of carbon stocks positively correlates with CO<sub>2</sub> sequestration.

The species with the lowest potential for CO<sub>2</sub> sequestration was *A. indica* due to the small tree number, young plant age, and small leaves that cause the process of CO<sub>2</sub> assimilation to be slightly hampered and tends to be ineffective. The amount of CO<sub>2</sub> the plant can absorb is affected by several factors, such as plant species, site conditions, and plant density. These factors significantly affect the amount of CO<sub>2</sub> sequestration by certain plants, impacting differences between plants. According to Zulkifli (2010), forests consisting of trees in the growth phase can absorb more CO<sub>2</sub>, while mature forests with small growth can store carbon stock, but the ability to absorb excess CO<sub>2</sub> is limited. On the other hand, forests can absorb carbon for a long time by implementing sustainable forest management. Therefore, planting vegetation on vacant land or rehabilitating damaged forests will help absorb excess CO<sub>2</sub> in the atmosphere.

## 4. Conclusions

This study identified 16 tree species in the BNI Urban Forest Banda Aceh, consisting of 8 families with a total of 318 individuals. *Casuarina equisetifolia* was the most common type of tree were 163 trees (51.3%). The results revealed that the above-ground biomass, carbon stock, and carbon sequestration by the vegetation in the BNI Urban Forest were 24.66 tons/ha, 11.59 tons/ha, and 42.51 tons/ha, respectively. This study provided data related to the existing conditions of the vegetation and the potential of carbon sequestration, which can be used to monitor and evaluate the benefits of urban forests in Banda Aceh.

## References

- Alfida, A., Hanum, U., and Eliyanti, E. 2016. Kupu-Kupu (Rhopalocera) di Kawasan Hutan Kota BNI Banda Aceh. *BIOTIK: Jurnal Ilmiah Biologi Teknologi dan Kependidikan* 4(2): 117. DOI: [10.22373/biotik.v4i2.2906](https://doi.org/10.22373/biotik.v4i2.2906)
- Ali, S., Khan, S. M., Siddiq, Z., Ahmad, Z., Ahmad, K. S., Abdullah, A., Hashem, A., Al-Arjani, A. B. F., and AbdAllah, E. F. 2022. Carbon Sequestration Potential of Reserve Forests Present in the Protected Margalla Hills National Park. *Journal of King Saud University* -

- Science* 34(4): 101978. DOI: [10.1016/j.jksus.2022.101978](https://doi.org/10.1016/j.jksus.2022.101978)
- Badan Standardisasi Nasional. 2011. Pengukuran dan Penghitungan Cadangan Karbon – Pengukuran Lapangan untuk Penaksiran Cadangan Karbon Hutan (Ground Based Forest Carbon Accounting). Indonesia.
- Baruna, T., Astiani, D., and Darwati, H. 2017. Estimation of Carbon Stock in Urban Forest of Pontianak City Hall. *Jurnal Hutan Lestari* 5: 547-555.
- Barona, O. C., Wolf, K., Kowalski, J. M., Kendal, D., Byrne, J. A., and Conway, T. M. 2022. Diversity in Public Perceptions of Urban Forests and Urban Trees: A Critical Review. *Landscape and Urban Planning* 226: 104466. DOI: [10.1016/j.landurbplan.2022.104466](https://doi.org/10.1016/j.landurbplan.2022.104466)
- Conway, T. M., and Vecht, J. V. 2015. Growing a Diverse Urban Forest: Species Selection Decisions by Practitioners Planting and Supplying Trees. *Landscape and Urban Planning* 138: 1-10. DOI: [10.1016/j.landurbplan.2015.01.007](https://doi.org/10.1016/j.landurbplan.2015.01.007)
- De Groot, M., Diaci, J., and Ogris, N. 2019. Forest Management History is an Important Factor in Bark Beetle Outbreaks: Lessons for the Future. *Forest Ecology and Management* 433: 467–474. DOI: [10.1016/j.foreco.2018.11.025](https://doi.org/10.1016/j.foreco.2018.11.025)
- Devisscher, T., Ordóñez-Barona, C., Dobbs, C., Dias Baptista, M., Navarro, N. M., Orozco Aguilar, L. A., Cercas Perez, J. F., Rojas Mancebo, Y., and Escobedo, F. J. 2022. Urban Forest Management and Governance in Latin America and the Caribbean: A Baseline Study of Stakeholder Views. *Urban Forestry and Urban Greening* 67: 127441. DOI: [10.1016/j.ufug.2021.127441](https://doi.org/10.1016/j.ufug.2021.127441)
- Fenton, P. 2017. The Role of Port Cities and Transnational Municipal Networks in Efforts to Reduce Greenhouse Gas Emissions on Land and at Sea from Shipping – An Assessment of the World Ports Climate Initiative. *Marine Policy* 75: 271-277. DOI: [10.1016/j.marpol.2015.12.012](https://doi.org/10.1016/j.marpol.2015.12.012)
- Harjadi, B. 2017. Peran Cemara Laut (*Casuarina equisetifolia*) dalam Perbaikan Iklim Mikro Lahan Pantai Berpasir di Kebumen. *Jurnal Penelitian Pengelolaan Daerah Aliran Sungai* 1(2): 73-81. DOI: [10.20886/jppdas.2017.1.2.73-81](https://doi.org/10.20886/jppdas.2017.1.2.73-81)
- Hyde, W. F., Olmos, V. M., Robalino, J., da Gama e Silva, Z. A. G. P., Susaeta, A., and Yin, R. 2022. Latin America: A Regional Perspective on Its Forest Policy and Economics. *Forest Policy and Economics* 141: 102760. DOI: [10.1016/j.forpol.2022.102760](https://doi.org/10.1016/j.forpol.2022.102760)
- Krisnawati, H., Adinugroho, W. C., and Imanuddin, R. 2012. *Model Model Alometrik untuk Pendugaan Biomassa Pohon pada Berbagai Tipe Ekosistem Hutan di Indonesia*. Monograf. Pusat Penelitian dan Pengembangan Konservasi dan Rehabilitasi. Badan Penelitian dan Pengembangan Kehutanan. Bogor, Indonesia. 65 p.
- Ledheng, L., Naisumu, Y. G., and Binsasi, R. 2022. The Estimation of Biomass in *Rhizophora apiculata* and *Rhizophora mucronata* in Tuamese Village, North Central Timor Regency, East Nusa Tenggara Province. *Jurnal Sylva Lestari* 10(1): 39-48. DOI: [10.23960/jsl.v10i1.555](https://doi.org/10.23960/jsl.v10i1.555)
- Lee, M. B., Zhang, M., Chan, S., Zhang, Q., Che, X., Wong, K. C., Zhao, X., and Zou, F. 2021. Environmental Determinants of Avian Species Richness in Forests in a Highly Urbanized Subtropical City: The Effects of Spatial Scale and Open Space Feature. *Global Ecology and Conservation* 30: e01802. DOI: [10.1016/j.gecco.2021.e01802](https://doi.org/10.1016/j.gecco.2021.e01802)
- Mahmoudi, B., Sorouri, Z., Zenner, E. K., and Mafi-Gholami, D. 2022. Development of a New Social Resilience Assessment Model for Urban Forest Parks. *Environmental Development* 43: 100724. DOI: [10.1016/j.envdev.2022.100724](https://doi.org/10.1016/j.envdev.2022.100724)

- Marchi, S., Black, B. A., Elkins-Tanton, L. T., and Bottke, W. F. 2016. Massive Impact-Induced Release of Carbon and Sulfur Gases in the Early Earth's Atmosphere. *Earth and Planetary Science Letters* 449: 96-104. DOI: [10.1016/j.epsl.2016.05.032](https://doi.org/10.1016/j.epsl.2016.05.032)
- Matthews, K. B., Wardell-Johnson, D., Miller, D., Fitton, N., Jones, E., Bathgate, S., Randle, T., Matthews, R., Smith, P., and Perks, M. 2020. Not Seeing the Carbon for the Trees? Why Area-Based Targets for Establishing New Woodlands Can Limit or Underplay Their Climate Change Mitigation Benefits. *Land Use Policy* 97: 104690. DOI: [10.1016/j.landusepol.2020.104690](https://doi.org/10.1016/j.landusepol.2020.104690)
- Morakinyo, T. E., Ouyang, W., Lau, K. K. L., Ren, C., and Ng, E. 2020. Right Tree, Right Place (Urban Canyon): Tree Species Selection Approach for Optimum Urban Heat Mitigation - Development and Evaluation. *Science of the Total Environment* 719: 137461. DOI: [10.1016/j.scitotenv.2020.137461](https://doi.org/10.1016/j.scitotenv.2020.137461)
- Mukhlison. 2013. Pemilihan Jenis Pohon untuk Pengembangan Hutan Kota di Kawasan Perkotaan Yogyakarta. *Jurnal Ilmu Kehutanan* 10(1): 37-47.
- Muslih, A. M., Bintoro, A., Duryat, Subhan, Arlita, T., Rasyid, U. H. A., Umam, A. H., Farida, A., and Yanti, L. A. 2021. The Study of Silvicultural Aspects of Community Forest in Buana Sakti Lampung Province. *IOP Conference Series: Earth and Environmental Science* 667(1): 12053. DOI: [10.1088/1755-1315/667/1/012053](https://doi.org/10.1088/1755-1315/667/1/012053)
- Muslih, A. M., Manalu, Y. H. B., Arlita, T., Moulana, R., and Anhar, A. 2022. Inventory of *Dryobalanops aromatica* Gaertn. in Lae Kombih Forest Park Area, Subulussalam. *IOP Conference Series: Earth and Environmental Science* 951(1): 12089. DOI: [10.1088/1755-1315/951/1/012089](https://doi.org/10.1088/1755-1315/951/1/012089)
- Ng, Y., Astiani, D., and Ekamawanti, H. A. 2021. Estimation of Tree Carbon Stocks in the Green Open Space of the Faculty of Forestry, Tanjungpura University. *Jurnal Sylva Lestari* 9(3): 434-443. DOI: [10.23960/jsl.v9i3.517](https://doi.org/10.23960/jsl.v9i3.517)
- Petri, A. C., Wilson, B., and Koeser, A. 2019. Planning the Urban Forest: Adding Microclimate Simulation to the Planner's Toolkit. *Land Use Policy* 88: 104117. DOI: [10.1016/j.landusepol.2019.104117](https://doi.org/10.1016/j.landusepol.2019.104117)
- Rahmadwiati, R., Wicaksono, R. L., and Apriyanto, D. 2022. Estimation and Economic Valuation of Above-Ground Carbon in Forest Area with the Special Purpose of Gunung Bromo, Karanganyar, Central Java, Indonesia. *Jurnal Sylva Lestari* 10(1): 116-126. DOI: [10.23960/jsl.v10i1.548](https://doi.org/10.23960/jsl.v10i1.548)
- Sharkey, T. D. 2021. *Photosynthesis | Photosynthetic Carbon Dioxide Fixation*. U.S. Department of Energy Office of Scientific and Technical Information. DOI: [10.1016/B978-0-12-819460-7.00011-6](https://doi.org/10.1016/B978-0-12-819460-7.00011-6)
- Skytt, T., Nielsen, S. N., and Jonsson, B.-G. 2020. Global Warming Potential and Absolute Global Temperature Change Potential from Carbon Dioxide and Methane Fluxes as Indicators of Regional Sustainability – A Case Study of Jämtland, Sweden. *Ecological Indicators* 110: 105831. DOI: [10.1016/j.ecolind.2019.105831](https://doi.org/10.1016/j.ecolind.2019.105831)
- Subhan, Umam, A. H., Muslih, A. M., Rasyid, U. H. A., Farida, A., and Anhar, A. 2021. Strategy Involving Stakeholders and Government to Contribute and to Implement the ILEDSA. *IOP Conference Series: Earth and Environmental Science* 711(1): 12029. DOI: [10.1088/1755-1315/711/1/012029](https://doi.org/10.1088/1755-1315/711/1/012029)
- Subhan, Anhar, A., Muslih, A. M., Ar-Rasyid, U. H., Maimunah, S., and Nasution, A. A. 2022. Urban Forest Carbon Stock and Biodiversity Assessment at Nagan Raya Regency. *IOP*

- Conference Series: Earth and Environmental Science* 951(1): 12071. DOI: [10.1088/1755-1315/951/1/012071](https://doi.org/10.1088/1755-1315/951/1/012071)
- Sugianto, S., Muslih, A. M., Ar-Rasyid, U. H., and Anhar, A. 2021. Vegetation Analysis of Rawa Singkil Wildlife Reserve in Rantau Gedang Village, Singkil Distric, Aceh Singkil Regency, Aceh Province. *IOP Conference Series: Earth and Environmental Science* 667(1): 12068. DOI: [10.1088/1755-1315/667/1/012068](https://doi.org/10.1088/1755-1315/667/1/012068)
- Sumardi, and Widyastuti, S. M. 2013. *Dasar-Dasar Perlindungan Hutan*. Gadjah Mada University Press. Yogyakarta.
- Sun, X., Li, J., Cameron, D., and Moore, G. 2022. Sap Flow Monitoring of Two Australian Native Tree Species in a Suburban Setting: Implications for Tree Selection and Management. *Plant Science* 317: 111194. DOI: [10.1016/j.plantsci.2022.111194](https://doi.org/10.1016/j.plantsci.2022.111194)
- Toochi, E. C. 2018. Carbon Sequestration: How Much Can Forestry Sequester CO<sub>2</sub>? *Forestry Research and Engineering: International Journal* 2(3): 148–150. DOI: [10.15406/freij.2018.02.00040](https://doi.org/10.15406/freij.2018.02.00040)
- Wahanisa, R. 2015. Model Pengelolaan Hutan Bersama Masyarakat (PHBM). *Yustisia Jurnal Hukum* 4(2): 416-438.
- Wu, J., Long, J., Liu, H., Sun, G., Li, J., Xu, L., and Xu, C. 2021. Biogenic Volatile Organic Compounds from 14 Landscape Woody Species: Tree Species Selection in the Construction of Urban Greenspace with Forest Healthcare Effects. *Journal of Environmental Management* 300: 113761. DOI: [10.1016/j.jenvman.2021.113761](https://doi.org/10.1016/j.jenvman.2021.113761)
- Xu, Y., and Cui, G. 2021. Influence of Spectral Characteristics of the Earth's Surface Radiation on the Greenhouse Effect: Principles and Mechanisms. *Atmospheric Environment* 244: 117908. DOI: [10.1016/j.atmosenv.2020.117908](https://doi.org/10.1016/j.atmosenv.2020.117908)
- Zulkifli, H. 2010. Pendidikan Lingkungan bagi Masyarakat sebagai Mitigasi Dampak Perubahan Iklim Melalui Upaya Penyimpanan Karbon pada Kawasan Hijau. *Majalah Ilmiah FORUM MIPA* 13(2): 111-116.